Biological Times

Volume No. 4 | Issue No. 10 | Pages 12-13

Beyond the Hype: Scaling Black Soldier Fly for Green Economy Pathways—Challenges and Innovations

Esha Nadeem[1*], Ali Raza[2], Hassan Raza[3] and Muhammad Hasnain[2]

1. Department of Zoology, Wildlife and Fisheries, University of Agriculture Faisalabad, Pakistan
2. Department of Entomology, University of Agriculture Faisalabad, Pakistan
3. Institute of Food Science and Technology, Khawaja Fareed University of Engineering & Information Technology (KFUEIT), Pakistan

*Corresponding Author: **eshanadeemofficial@gmail.com**

ABSTRACT

The Black Soldier Fly (BSF), Hermetia illucens, has rightfully emerged as a flagship species in the pursuit of a circular bioeconomy. Its larvae offer a triple-value proposition: rapid organic waste reduction, a sustainable source of protein and lipids for animal feed, and the production of a nutrient-rich organic fertilizer (frass). While extensive research has validated its efficacy at a laboratory and small-scale level, the transition to widespread industrial adoption presents a complex array of scientific, economic, and regulatory challenges. This review extension article moves beyond the established baseline to critically examine the key hurdles hindering the sector's maturation. We identify and discuss five critical frontiers: (1) the safety and regulatory landscape concerning substrate contaminants and product standardization; (2) the technological and economic bottlenecks in industrial-scale production; (3) the nutritional optimization and value-added applications of BSF products; (4) the environmental and social impact assessments required for credible lifecycle analysis; and (5) the policy frameworks and consumer acceptance needed for market integration.Advancing BSF research requires interdisciplinary efforts and collaboration among policymakers, scientists, and industry. The coming decade is likely to determine whether BSF transitions from pilot projects to mainstream solutions.

Keywords: *Hermetia illucens*, circular bioeconomy, waste valorization, industrial automation, regulatory policy, consumer acceptance

To cite this article: Nadeem E, A Raza, H Raza & M Hasnain. Beyond the Hype: Scaling Black Soldier Fly for Green Economy Pathways—Challenges and Innovations. Biological Times. 2025. October 4(10): 12-13.

Introduction

The escalating crisis of organic waste generation, combined with the environmental burden of conventional protein sources like fishmeal and soybean meal, has driven the search for innovative solutions. Among these, the Black Soldier Fly (BSF) stands out due to its efficiency in bioconversion. Foundational studies have shown that BSF larvae can consume diverse organic substrates, reducing waste volume by 50–70% within a short period [1]. Their biomass is rich in protein (35–45%) and lipids (20–30%), offering a sustainable alternative for aquaculture and poultry feed [2, 3]. The frass, rich in nitrogen, phosphorus, and potassium, completes the cycle as an effective organic fertilizer [4]. While these benefits are well-demonstrated at pilot scales, significant gaps remain between proof-of-concept and industrially viable implementation. The BSF industry stands at a critical juncture where addressing safety, regulation, technology, nutrition, and socio-economic integration will define its long-term impact. This review provides a critical analysis of research and development priorities, arguing that a holistic, interdisciplinary approach is essential for embedding BSF technology in the global circular economy.

Critical Frontier I: Ensuring Safety and Navigating the Regulatory Labyrinth

The strength of BSF lies in its ability to thrive on diverse organic waste streams, but this raises serious concerns regarding safety and regulation.

The Contaminant Question

A major challenge is the potential bioaccumulation of contaminants in larval biomass and frass. While BSF larvae reduce pathogen loads due to their antimicrobial properties [5], they can accumulate heavy metals such as cadmium, lead, and arsenic, depending on the substrate [6]. This creates risks if contaminated larvae enter the food chain. Similarly, the fate of mycotoxins, pesticides, pharmaceutical residues, and microplastics in BSF systems remains insufficiently understood [7]. To address these gaps, future research must prioritize comprehensive risk profiling by mapping contaminant transfer factors across various waste streams, elucidating detoxification mechanisms employed by larvae, and developing pre-treatment standards for high-risk substrates.

The Need for Harmonized Regulations

Globally, BSF regulations remain fragmented and inconsistent. The European Union, under Regulation (EU) 2017/893, permits BSF protein in aquaculture and poultry feed but restricts substrate choices [8]. Many other countries either lack clear regulations or impose overly restrictive policies that limit industry growth. Key needs include internationally standardized product specifications for BSF meal, oil, and frass to ensure nutritional consistency and safety; science-based frameworks for approving feed substrates instead of narrow "positive lists"; and formal certification criteria for frass as organic fertilizer, focusing on stability, maturity, and contaminant safety.

Critical Frontier II: Technological Innovation and Economic Viability at Scale

Transitioning BSF production from artisanal setups to industrial levels requires overcoming efficiency bottlenecks through automation and rigorous sustainability assessments.

The Automation Imperative

Current production practices rely heavily on manual labor for egg collection, larval sorting, and frass separation, limiting economic scalability. Industrial viability depends on automation. Smart rearing systems using sensor-based monitoring of temperature, humidity, and larval biomass, integrated with automated climate control, can optimize conditions and reduce mortality. Robotics for harvesting and processing—such as mechanical systems to separate prepupae from frass and automated systems for drying, grinding, and oil extraction—would improve yield and consistency. Data-driven optimization through AI and machine learning offers further opportunities to predict yields, refine feeding regimes, and reduce inefficiencies.

Rigorous Life Cycle Assessment (LCA)

Although BSF is promoted as a green technology, industrial-scale operations must undergo full life cycle assessments (LCAs) to validate sustainability claims. Energy-intensive processes such as climate control, feed grinding, and larval drying could undermine benefits if powered by fossil fuels. Assessing the net energy balance is therefore critical. Similarly, water use for larval humidity maintenance and facility cleaning must be quantified. Standardized LCA methodologies should compare BSF not only with landfilling but also with other valorization methods such as anaerobic digestion and composting, using consistent functional units such as "treatment of one metric ton of organic waste."

Critical Frontier III: Nutritional Optimization and Value-Added Applications

Maximizing the commercial potential of BSF requires nutritional tailoring and exploration of high-value markets beyond aquaculture and poultry

Tailoring Larval Nutrition

The nutritional profile of BSF larvae reflects their diet [10], offering opportunities for precision nutrition. Substrate manipulation can enrich larvae with specific fatty acids such as omega-3s by incorporating marine algae residues. Biofortification approaches could increase micronutrient levels, producing functional feed ingredients with added value for animal health. Additionally, BSF oil, rich in lauric acid and other bioactive

compounds, holds potential as a natural antimicrobial alternative to antibiotics.

Expanding the Application Horizon

Beyond aquaculture and poultry, other sectors show promise. In monogastric livestock, BSF meal could support gut health in pigs, while in the pet food industry, it is valued as a novel, high-protein ingredient for premium diets. The larval exoskeleton provides chitin, a versatile biopolymer with applications in pharmaceuticals, cosmetics, and biodegradable plastics [11]. Developing efficient chitin extraction techniques could unlock new revenue streams, diversifying the BSF product portfolio.

Critical Frontier IV: Integrating Social and Environmental Impact Assessments

The industry's success will depend not only on technological and economic viability but also on equitable societal integration and ecological safeguards.

Social Equity and Inclusivity

BSF farming has potential to generate green jobs, especially in developing countries with acute waste challenges. Ensuring inclusivity is crucial to prevent monopolization by large corporations. Smallholder integration through accessible technologies and business models should be prioritized. Gender inclusivity is also important, given women's significant role in waste management and small-scale livestock production [12]. Knowledge transfer via open-access platforms and extension services will be essential to democratize innovation.

Biodiversity and Ecosystem Considerations

Scaling BSF production requires evaluating potential ecological impacts. Although BSF is generally non-pestilent, accidental escape into non-native ecosystems could disrupt local insect communities and decomposition dynamics [13]. Therefore, robust containment strategies and environmental risk assessments are necessary components of industrial planning.

Critical Frontier V: Policy Frameworks and Market Creation

Technological progress alone cannot secure the industry's future; enabling policy and market mechanisms are equally critical.

Supportive Policy Instruments

Governments can accelerate BSF adoption through policies that reflect the true environmental costs of waste disposal and conventional feed production. Tax incentives and subsidies could encourage BSF investments, while Extended Producer Responsibility (EPR) frameworks would ensure consistent waste feedstock supplies by holding food producers accountable for organic waste management. Public procurement policies that favor products raised on sustainable feed ingredients, such as BSF meal, could further stimulate demand.

The Consumer Acceptance Challenge

Consumer acceptance remains a key determinant of market growth. Although psychological aversion is less pronounced in animal feed compared to direct human diets, it still shapes perceptions of insect-fed products like poultry or fish [14]. Transparent communication of safety and sustainability benefits, backed by strong regulatory assurances, will be essential for building trust and securing long-term adoption.

Conclusion and Synthesis

The Black Soldier Fly presents a paradigm-shifting opportunity to address some of the most pressing challenges of the 21st century: waste management, food security, and climate change. The foundational research, as summarized in the initial review, provides an unequivocal proof-of-concept. However, this extended analysis demonstrates that the path forward is complex and requires a concerted, multi-stakeholder effort.

The five frontiers outlined—safety and regulation, scale-up economics, product valorization, socio-environmental integration, and policy support—are deeply interconnected. Progress in automating production (Frontier II) will impact economic viability and LCA outcomes. Advances in understanding contaminant transfer (Frontier I) are prerequisites for robust regulatory frameworks (Frontier V). The industry's growth will therefore depend on collaboration between entomologists, chemical engineers, economists, social scientists, policymakers, and industry partners.

The next decade will be decisive. By channeling research investments and policy support towards these critical areas, we can transition Black Soldier Fly bioconversion from a promising circular economy prototype into a scalable, safe, and sustainable industry. In doing so, we can move closer to a future where waste is not an endpoint but a beginning, and where resource cycles are not linear but beautifully, and productively, circular.

References

[1] Diener S, Solano NMS, Gutiérrez FR, Zurbrügg C, Tockner K. Biological treatment of municipal organic waste using black soldier fly larvae. Waste Biomass Valor. 2011;2(4):357-63.

[2] 2. Henry M, Gasco L, Piccolo G, Fountoulaki E. Review on the use of insects in the diet of farmed fish: Past and future. Anim Feed Sci Technol. 2015;203:1-22.

[3] 3. Biasato I, Gasco L, De Marco M, Renna M, Rotolo L, Dabbou S, et al. Yellow mealworm larvae (Tenebrio molitor) inclusion in diets for male broiler chickens: Effects on growth performance, gut morphology, and histological findings. Poult Sci. 2018;97(2):540-8.

[4] 4. Beesigamukama D, Mochoge B, Korir N, Fiaboe KK, Nakimbugwe D, Khamis FM. Exploring black soldier fly frass as novel fertilizer for improved growth, yield, and nitrogen use efficiency of maize. Front Plant Sci. 2020;11:574592.

[5] 5. Erickson MC, Islam M, Sheppard C, Liao J, Doyle MP. Reduction of Escherichia coli O157:H7 and Salmonella enterica serovar Enteritidis in chicken manure by larvae of the black soldier fly. J Food Prot. 2004;67(4):685-90.

[6] 6. Diener S, Zurbrügg C, Tockner K. Bioaccumulation of heavy metals in the black soldier fly, Hermetia illucens and effects on its life cycle. J Insects Food Feed. 2015;1(4):261-70.

[7] 7. van der Fels-Klerx HJ, Camenzuli L, van der Lee MK, Oonincx DGAB. Uptake of cadmium, lead and arsenic by Tenebrio molitor and Hermetia illucens from contaminated substrates. PLoS One. 2018;13(11):e0207512.

[8] 8. European Commission. Commission Regulation (EU) 2017/893 of 24 May 2017 amending Annexes I and IV to Regulation (EC) No 999/2001 and Annexes X, XIV and XV to Commission Regulation (EU) No 142/2011 as regards processed animal protein. Off J Eur Union. 2017;L138:92.

[9] 9. Smetana S, Schmitt E, Mathys A. Sustainable use of Hermetia illucens insect biomass for feed and food: Attributional and consequential life cycle assessment. Resour Conserv Recycl. 2019;144:285-96.

[10] 10. Spranghers T, Ottoboni M, Klootwijk C, Ovyn A, Deboosere S, De Meulenaer B, et al. Nutritional composition of black soldier fly (Hermetia illucens) larvae as affected by the feed substrate. J Insects Food Feed. 2017;3(2):105-20.

[11] 11. Kaya M, Baran T, Karaarslan M. A new method for fast chitin extraction from shells of crab, crayfish and shrimp. Nat Prod Res. 2015;29(15):1477-80.

[12] 12. Nyakeri E, Ayieko M, Amimo F. An open system for farming black soldier fly larvae as a source of proteins for smallscale poultry and fish production. J Insects Food Feed. 2017;3(1):51-6.

[13] 13. Bosch G, Van Zanten HH, Zamprogna A, Veenenbos M, Meijer NP, Van der Fels-Klerx HJ, et al. Conversion of organic resources by black soldier fly larvae: Legislation, efficiency and environmental impact. J Clean Prod. 2019;222:355-63.

[14] 14. Verbeke W, Spranghers T, De Clercq P, De Smet S, Sas B, Eeckhout M. Insects in animal feed: Acceptance and its determinants among farmers, agriculture sector stakeholders and citizens. Anim Feed Sci Technol. 2015;204:72-87.

[15] 15. Wang YS, Shelomi M. Review of black soldier fly (Hermetia illucens) as animal feed and human food. Foods. 2017;6(10):91.

[16] 16. Gold M, Tomberlin JK, Diener S, Zurbrügg C, Mathys A. Decomposition of biowaste macronutrients, microbes, and chemicals in black soldier fly larval treatment: A review. Waste Manag. 2018;82:302-18.

[17] 17. Barragan-Fonseca KB, Dicke M, van Loon JJ. Nutritional value of the black soldier fly (Hermetia illucens L.) and its suitability as animal feed. J Insects Food Feed. 2017;3(2):105-20.

[18] 18. Lalander C, Diener S, Magri ME, Zurbrügg C, Lindström A, Vinnerås B. Faecal sludge management with the larvae of the black soldier fly (Hermetia illucens)—From a hygiene aspect. Sci Total Environ. 2013;458:312-18.

[19] 19. Sheppard DC, Newton GL, Thompson SA, Savage S. A value added manure management system using the black soldier fly. Bioresour Technol. 1994;50(3):275-9.

[20] 20. Čičková H, Newton GL, Lacy RC, Kozánek M. The use of fly larvae for organic waste treatment. Waste Manag. 2015;35:68-80.