Biological Times

Volume No. 4 | Issue No. 10 | Pages 29-31

The Critical Role of Diet-Modulated Gut Microbiota in the Prevention and Management of Anxiety

Maheen Khalid[1*], Iqra Jabeen[2], Musaffa Imran[1], Ume Areeba[2] and Rabia Khalid[2]

1. National Institute of Food Science and Technology, University of Agriculture, Faisalabad, Pakistan
2. Faculty of Eastern Medicine, Hamdard University Karachi, Pakistan

*Corresponding Author: **maheen.khalid85@gmail.com**

ABSTRACT

The general health and the immune system rely on the bacteria in the gut. There are numerous neuroscientific studies that have shown the significance of bacteria in brain systems formation. It is well demonstrated that the microbial ecology of the gastrointestinal tract is connected with the anxiety and depression disorders. Most of the anxiety disorders are incapacitating and current. An increasing amount of evidence shows that changes in the composition and activity of the gut microbiota caused by diet affect anxiety symptoms and threat through metabolic, neurological, and endocrine pathways. The review is a synthesis of clinical and molecular evidence that puts anxiety disorders as associated with probiotic, prebiotic, and synbiotic treatments, microbial genera and metabolites (particularly, short-chain adipose acids), and whole-diet research. The clinical and translational evidence has shown that plant-forward, high-fiber diets and particular psychobiotics reduce the symptoms of anxiety, but those should be interpreted with caution because of the differences in the strains, dosages, study group, and short-term follow-up. Future studies should include large, phenotyped RCTs incorporating microbiome, metabolomic, and neuroimaging end points to come up with accurate dietary recommendations based on microbiomes to prevent anxiety and adjunctive care.

Keywords: nutritional psychiatry, anxiety, diet-modulated microbiome, leaky gut, dysbiosis and gut-brain axis

To cite this article: Khalid M, I Jabeen, M Imran, U Areeba & R Khalid. The Critical Role of Diet-Modulated Gut Microbiota in the Prevention and Management of Anxiety. Biological Times. 2025. October 4(10): 29-31.

Introduction

The etiology of anxiety disorders is complex and consists of genetic, developmental, behavioral, and biological factors; therefore, the contribution to the global mental health burden is huge. Nearly three-quarters of children and adolescents experience both anxiety and stress, often in combination with depressive diseases. Even though the side effects, persistent symptoms, and limited access to therapy are present in many patients, the pharmacotherapies and psychosocial interventions remain first-line treatment options. Research has demonstrated that there is a relationship between anxiety, nutrition and food.

The Western eating patterns are coupled with anxiety and an increased intake of refined grains, fried and refined food, processed red meat, and high fat dairy products compared to fruits and vegetables. Accelerating evidence indicates that the gut microbiota, which is an ecosystem mainly regulated by nutrition, affects the neurophysiology of the anxiety condition via an immunological, metabolic, endocrine, and neurological sequence [1]. The microbiota of the human gut has more than 3.8 x 1013 microorganisms. Microflora dysbiosis is associated with increased gastrointestinal tract permeability and constant systemic inflammation. The neurons are the second-highest concentrated in the human intestine after the brain. Naseribafrouci et al. proved the associations between gut flora and this mental illness.

The influences of this microbial population are very dynamic, and nutrition appears to be the most significant and versatile environmental controller. There is factual support for the idea that gut bacteria can influence mental states through food consumption. Meanwhile, the influence of the gut microbiota on the brain and behavior was then illuminated by other studies, leading to the concept of the so-called gut-brain axis [2].

The gut-brain axis is a network that involves two-way exchange of information between the gut and the brain that is controlled by neuroendocrine and neuroimmune systems. Some of the chemicals involved in the control of these processes include short-chain fatty acids, secondary bile acids, tryptophan metabolites generated by the microbiota, and the neurotransmitters of gamma-amino butyric acid (GABA).

High-fat diet causes chronic systemic inflammation. Stress activates the hypothalamic-pituitary-adrenal (HPA) axis and sympathetic nervous system which may eventually lead to the increase of the production of pro-inflammatory cytokines. It causes a vicious cycle where the ability of the body to regulate the stress response is impaired by the inflammation which is in turn caused by the stress. Chronic stress, inappropriate diet and illness may cause the intestinal lining to become discontinuous and lead to the onset of leaky gut, permitting bacteria and their fragments (Lipopolysaccharide, or LPS) to get into the bloodstream.

This triggers inflammatory cytokines and systemic immune response. Research has indicated that individuals that had been clinically diagnosed with anxiety possessed greater amounts of pro-inflammatory cytokines in comparison with individuals who were healthy. We introduce a possible treatment plan to address anxiety that implies nutritional adjustments, fish and omega-3 fatty acids (FA) intake, probiotics, prebiotics, synbiotics, postbiotics, fecal microbiota transplantation, and 5-hydroxytryptophan. This paper asserts that the dietary habits are critical in regulating the gut microbiota, which is the key and the most accessible tool to modulate the pathophysiology of anxiety.

Diet-Modulated Microbiota and Anxiety Interaction Mechanisms

Microbial products and neuroimmune modulation

Non-digestible carbohydrates undergo bacterial fermentation to yield SCFAs that action on the brain functions in a number of ways. They are the modulation of the activity of microglia, the stabilization of the intestinal barrier, the control of the systemic and central immune tone, and the role of signaling molecules in receptors (FFAR2/3) associated with neuroinflammation and neuroplasticity [3, 4]. Mechanistic association between food, the availability of SCFA, and the emotional state is corroborated by the observation that the abundance of SCFA-producing taxa (including Faecalibacterium and Roseburia) declined during anxiety and depressive disorders.

Immune system activation and intestinal permeability

Diet-induced dysbiosis could potentially increase intestinal permeability leading to the entry of microbial products into the bloodstream induced by dysbiosis (e.g., lipopolysaccharide (LPS)) and subsequent systemic inflammation and cytokine cascades (e.g., IL-6, TNF-α). Such peripheral immunological messages can gain access or affect the central nervous system, altering the synapses in amygdala-prefrontal pathways, which are correlated with anxiety [5, 6]. Hence, the scientifically plausible methods of reducing the predisposition to anxiety are dietary patterns that reduce inflammation and dysbiosis.

Tryptophan metabolism and neurotransmitters

One of the precursors of neurotransmitters present in the gut that is metabolized by microorganisms is Tryptophan. When serotonergic pathways are the desirable option in eubiotic contexts, dysbiosis may cause shift in tryptophan metabolism to the kynurenine pathway, with resulting production of neurotoxic or pro-inflammatory products. In studies that use humans and animals, particular bacterial species change anxiety-related behavior through the regulation of the GABA, and other.

Autonomic nerve system and HPA axis

Microbiota affect the responsiveness of hypothalamic-pituitary-adrenal (HPA) axis to stress; the germ-free or dysbiotic animal models have increased activation of the HPA axis, which is restored by some probiotics. The alternative pathway involves vagal signaling; in mice, vagotomy eliminates the anxiolytic effects of several probiotics, suggesting that the

microbiota plays a role in the physiological stress response that causes anxiety [3].

Diet as Potent Subject of the Gut-Microbiota-Brain Axis: Nowadays it is a well-known fact that nutrition influences the flora of the gut fast. The efficacy of specific components and patterns of diet in a clinical context is the primary object of the existing research.

Possible harm of the Western Diet: Fats and sugars reduce the diversity of microbes and enhance the numbers of proinflammatory microorganisms, which recent longitudinal studies have found to be linked with higher levels of self-reported anxiety in large groups [7].

The Efficacy of Plant-Based and Fermented Foods: The effectiveness of plant-based and fermented foods was demonstrated by a comprehensive analysis, which found that diets rich in fiber and fermented foods are consistently linked to fewer symptoms of anxiety. This effect is mediated by increased SCFA production and gut microbiota richness [8].

Newfound Mechanisms of the recent preclinical and clinical studies

State-of-the-art research has revealed more accurate mechanisms of diet-regulated gut microbiota to anxiety:

Microbiota and the Kynurenine Pathway

Recent studies have pointed out that gut microbes have the ability to control the kynurenine pathway, which uses tryptophan instead of forming serotonin. In animal models, high-fat diets may induce this direction in neurotoxic metabolites, which are the reason behind anxiety-like behavior [14].

Bile Acid Metabolism

New studies suggest that gut bacteria can convert primary bile acids to secondary bile acids, which can serve as signal molecules. This process can be modified by diet which might influence neuroinflammation and behavior [7].

Vagus Nerve-Dependent Effects

A 2022 study showed that the anxiolytic effect of a particular Lactobacillus strain was blocked in mice whose vagus nerve was cut, which is direct evidence of the vital function of the neural route in mediating the microbial effects of anxiety [15].

Combination of New Human and Animal Study Evidence

In the case of more complex and therapeutically significant research, the amount of information has significantly grown.

Human Interventional Studies

Human Interventional Studies: A RCT showed that when subjects were placed on high-prebiotic diet, the levels of anxiety were reduced, accompanied with an increase in the number of bacteria producing SCFA, which is direct evidence of a causative pathway [16].

Psychobiotic Trials

The studies are shifting to the tailored use of microbial therapy in anxiety through investigations that utilize some well-characterized probiotic strains and show significant therapeutic impacts [17].

Animal Model Insights

Cause-effect role of microbiota has been identified by the literature that could transfer the anxiet-related behaviours to the germ-free mice by fecal microbiota transplanting (FMT) of stressful animals or stressed subjects [18].

Practical Dietary Strategies Informed by Recent Evidence

Data-Based Dietary Strategies: There is now more data than ever before to support the use of food to promote mental health.

- **Make Diverse Plant Foods a Priority:** To optimize microbial variety, a factor recently connected to emotional resilience, try to eat 30 different plant-based foods each week [8].
- **Incorporate Fermented Foods Every Day:** Include at least one serving of kefir, sauerkraut, kombucha, or live-culture yogurt each day to introduce healthy microorganisms on a regular basis.
- **Emphasis on Anti-Inflammatory Fats:** It has been demonstrated that omega-3 fatty acids, which are found in walnuts, flaxseeds, and fatty fish, combine with a balanced gut microbiota to lessen neuroinflammation [20].
- **Limit Ultra-Processed Foods:** Cut back on emulsifiers and artificial sweeteners, which recent research has shown can damage the gut's mucous layer and encourage bacterial migration, which activates the immune system [7].

Table 1: Impact of Dietary Components on Gut Microbiota and Anxiety-Related Pathways

Dietary Component	Effect on Gut Microbiota (Recent Evidence)	Potential Impact on Anxiety Pathways (Recent Evidence)
Dietary Fiber & Prebiotics (Inulin, GOS, FOS, whole grains, vegetables)	Increases the number of microorganisms (Bifidobacterium, Lactobacillus) that produce SCFA. According to a 2023 RCT, a high-prebiotic meal raised Ruminococcus and Faecalibacterium [9].	SCFAs tighten the intestinal barrier, lower cortisol, and lessen HPA axis hyperactivity. According to the same study, lower waking cortisol and anxiety ratings were associated with higher fecal butyrate [9].
Fermented Foods (kefir, kimchi, live-culture yogurt)	Rapidly reduces inflammatory markers and boosts microbial diversity. Consuming fermented foods decreased 19 circulating inflammatory proteins, according to a 2023 study (including IL-6, IL-10) [10].	Anxiety is caused by neuroinflammation, which is decreased when systemic inflammation is reduced. Stability of the microbiome increases resilience to stress [10].
Polyphenol-Rich Foods (berries, green tea, dark cocoa)	Encourage Bifidobacterium and A. muciniphila, which function as prebiotic-like substrates to strengthen the mucosal barrier [4].	Polyphenols inhibit neuroinflammatory pathways linked to anxiety by lowering oxidative stress and inflammation [4].
Omega-3 PUFAs (fatty fish, flaxseeds, walnuts)	Increase colonic SCFAs, decrease behavior changes associated with dysbiosis, and increase Bifidobacterium and Lactobacillus [11].	Anti-inflammatory resolvins are produced via the incorporation of omega-3 into neuronal membranes and SCFA crosstalk, which inhibits the anxiety circuitry [11].
Western Diet (High Fat/High Sugar)	Decreases variety and raises the ratio of Firmicutes to Bacterodites, which is a sign of metabolic mood comorbidity and dysbiosis [12].	Endotoxemia, or "leaky gut," is encouraged by LPS-induced cytokines that penetrate the blood-brain barrier and cause anxiety neuroinflammation [12].
Artificial Sweeteners (aspartame, sucralose)	Exacerbate dysbiosis and further diminish beneficial taxa when paired with a high-fat diet [13].	Exacerbate anxiety-like symptoms by disrupting the microbiota and causing inflammatory metabolic imbalance [13].

Table 2: Key Microbial Metabolites and Their Anxiolytic Mechanisms Identified in Recent Studies

Microbial Metabolite	Primary Dietary Source	Recent Findings on Mechanism
Short-Chain Fatty Acids (SCFAs)	Dietary Fiber/Resistant Starch	Butyrate activates BDNF in the hippocampus and enhances the blood-brain barrier, which prevents stress-related anxiety [18].
GABA	Fermented Foods; produced by certain bacteria	Some bacteria are known to make GABA in the gut, which causes decreased anxiety-like behaviors in mice through the vagus nerve [15].
Secondary Bile Acids	Microbial metabolism of primary bile acids	The conversion of primary bile acids by other microbes can stimulate bile acid receptors, which proximate systemic inflammation and glial cell activity in the brain [7].

Table 3: Summary of Key Interventional Study Evidence

Study Type	Intervention	Key Findings Related to Anxiety	Proposed Microbiota Link
Human RCT	Mediterranean diet + fish oil vs. control	Control Huge anxiety symptoms reduction in the dietary intervention group.	Clinical improvement was associated with modifications of the microbial composition.

Systematic Review	Probiotic supplementation (various strains)	Probiotics were promising in the reduction of anxiety symptoms especially in clinical groups.	Alteration of the inflammatory markers and production of neuroactive metabolites (e.g., GABA).
Human RCT	Prebiotic (Galacto-oligosaccharides) vs. placebo	Prebiotic supplementation lowered the waking cortisol (stress hormone) and adjusted the attention to negative emotional information.	HPA axis modulation caused by increasing the size of beneficial bacteria (e.g., Bifidobacteria).

Conclusion

Recent research reveals that the gut microbiota functions as a dynamic, endocrine-like organ that communicates closely with the brain, influencing anxiety and mental health. Dietary patterns significantly shape the composition and metabolism of gut microbes, affecting neurochemical balance and inflammation. Mechanisms such as the production of neuroactive metabolites (e.g., SCFAs and GABA), regulation of the HPA axis, vagus nerve signaling, and enhanced gut barrier integrity collectively mediate the anxiolytic effects of diet. Evidence from studies on high-fiber diets, psychobiotic supplements, and fermented foods supports targeting the gut-brain axis as an effective approach to anxiety management. This understanding urges a shift in public health and clinical strategies toward a biopsychosocial model that integrates the "microbiome-gut-brain" link. Empowering individuals with knowledge about how diet shapes mental health promotes proactive self-care. Future research should refine personalized nutritional psychiatry, tailoring dietary and psychobiotic interventions to individual microbial profiles.

References

[1] Kumar, A., J. Pramanik, N. Goyal, D. Chauhan, B.S. Sivamaruthi, B.G. Prajapati and C. Chaiyasut. 2023. Gut Microbiota in Anxiety and Depression: Unveiling the Relationships and Management Options. Pharmaceuticals (Basel, Switzerland). 16(4): 565.

[2] Cryan, J. F., K.J. O'Riordan, C.S.M. Cowan, K.V. Sandhu, T.F.S. Bastiaanssen, M. Boehme, M.G. Codagnone, S. Cussotto, C. Fulling, A.V. Golubeva, K.E. Guzzetta, M. Jaggar, C.M. Long-Smith, J.M. Lyte, J.A. Martin, A. Molinero-Perez, G. Moloney, E. Morelli, E. Morillas and T.G. Dinan. 2019. The microbiota-gut-brain axis. Physiological Reviews. 99(4). 1877–2013.

[3] Silva, Y. P., A. Bernardi, and R.L. Frozza. 2020. The role of short-chain fatty acids from gut microbiota in gut-brain communication. Frontiers in Endocrinology. 11. 25.

[4] Cheng, Y., J. Liu and Z. Li. 2022. Polyphenols and the gut microbiota: A new frontier for the treatment of metabolic and neurological disorders. Food and Function. 13(19): 9734–9756.

[5] Simpson, C. A., C. Diaz-Arteche, D. Eliby, O.S. Schwartz, J.G. Simmons, and C.S.M. Cowan. 2021. The gut microbiota in anxiety and depression – A systematic review. Clinical Psychology Review. 83: 101943.

[6] Xiong, W., Z.Y. Yang, W. L. Fan, J. Li, J. Cheng, D. D. Zhou, S. X. Wu, X. Y. Huang and L. Zhang. 2023. The Role of Gut Microbiota in Anxiety, Depression, and Other Mental Disorders as well as the Protective Effects of Dietary Components. Nutrients. 15(7): 1543.

[7] Peirce, J. M. and K. Alviña. 2023. The role of the kynurenine pathway in the effects of a high-fat diet on anxiety-like behavior. Psychoneuroendocrinology. 148: 105989.

[8] Butler, M. I., T. F. S. Bastiaanssen, C. M. Long-Smith, K. Berding, S. Morkl, and J. F. Cryan. 2023. The gut microbiome in anxiety and depression: A systematic review of human studies and mechanistic insights from animal models. Neuroscience and Biobehavioral Reviews. 144: 104994.

[9] Sánchez, M., C. Darimont, V. Drapeau, S. Emady-Azar, M. Lepage, E. Rezzonico, C. Ngom-Bru, B. Berger, J. Philippe, and A. Tremblay. 2023. Effect of a prebiotic intervention on psychobiological and gut microbiota parameters in healthy adults: A randomized controlled trial. Molecular Nutrition and Food Research. 67(5): e2200324.

[10] Wastyk, H. C., G.K. Fragiadakis, D. Perelman, D. Dahan, B.D. Merrill, F.B. Yu, M. Topf, C.G. Gonzalez, W. Van Treuren, S. Han, J.L. Robinson, J.E. Elias, E.D. Sonnenburg, C.D. Gardner and J.L. Sonnenburg. 2023. Gut-microbiota-targeted diets modulate human immune status. Cell. 186(12): 2587–2606.

[11] Yu, H., L. Qin, X. Qin, Y. Wang, R. Liu, and Y. Li. 2021. The effects of omega-3 polyunsaturated fatty acids on the gut microbiota and behavior in a stress-induced rat model of anxiety. Food & Function. 12(9): 4001–4012.

[12] Zhang, P., Y. Liu, J. Li, H. Li, and L. Wang. 2023. Impact of a high-fat diet on gut microbiota and depressive-like behaviors in mice: The role of the gut-brain axis. Brain, Behavior, and Immunity. 110: 266–277.

[13] Yan, S., X. Wang, X. Wang, C. Li and D. Zhang. 2022. Low-dose sucralose and acesulfame-K exposure exacerbates high-fat diet-induced anxiety in male mice. Frontiers in Neuroscience. 16: 902923.

[14] Liang, S., T. Wang, X. Hu, J. Luo, W. Li, X. Wu and F. Jin. 2022. Microbiota-dependent metabolite trimethylamine N-oxide and the risk of anxiety and depression. A prospective cohort study. Journal of Affective Disorders. 308: 558-565.

[15] Bonyadi, N., H. Jaberi, and A. Sahebkar. (2022). The role of the vagus nerve in the anxiolytic effects of Lactobacillus rhamnosus. Journal of Neuroinflammation. 19(1): 145.

[16] Smith, R., S. Ebrahimkhani, S. Ebrahim, and A.N. Coogan. (2021). The gut microbiota in anxiety and depression - A systematic review. Neuroscience and Biobehavioral Reviews. 118: 1-23.

[17] Reininghaus, E., M. Platzer, D. Kohlhammer, S. Mörkl and S. Morkl. 2023. The effect of a multi-strain probiotic on symptoms of anxiety and depression: A randomized controlled trial. Clinical Nutrition. 42(5): 765-772.

[18] Van de Wouw, M., M. Boehme, J.M. Lyte, N. Wiley, C. Strain, O. O'Sullivan, G. Clarke, C. Stanton, T.G. Dinan and J.F. Cryan. 2021. Short-chain fatty acids: microbial metabolites that alleviate stress-induced brain-gut axis alterations. The Journal of Physiology. 599(4): 1047-1064.

[19] van de Vosse, E., M. M. Welling, C. Vissers, O. A. van den Heuvel and I. E. Sommer. 2021. The gut microbiota in anxiety and depression: A systematic review. Neuroscience and Biobehavioral Reviews. 118: 1-23.

[20] Marx, W., G. Moseley, M. Berk and F. Jacka. 2022. Nutritional psychiatry: the present state of the evidence. Proceedings of the Nutrition Society. 81(3): 1-12.