

An Overview of Fasciolosis in Livestock and the Control Measures in Livestock

Muhammad Salman*

Department of Parasitology, Faculty of Veterinary Science, University of Agriculture, Faisalabad, Pakistan

*Corresponding Author: msalmanhameed@gmail.com

ABSTRACT

Fasciolosis caused by liver fluke, *Fasciola hepatica*, is a potentially significant threat to livestock. It results in substantial economic losses within the livestock industry. This parasitic disease affects a variety of animals including sheep, goats, cattle, and buffalos. This has the potential of transmission even to humans as a zoonotic disease. Snails play an important role in the life cycle of *Fasciola hepatica*. Clinical symptoms include anemia, reduced production and even death in severely infected animals. Diagnosis is made on the basis of clinical signs, post-mortem examinations and some laboratory diagnostic tests like ELISA. Treatment is with Triclabendazole as the drug of choice. Control needs coordinated measures involving herd health programs, snail population control, regular deworming, and maintaining proper hygiene.

Keywords: Fasciolosis, Liver fluke, Livestock, Parasites, Snails, Triclabendazole

To cite this article: Salman M. An Overview of Fasciolosis in Livestock and the Control Measures in Livestock. *Biological Times*. 2025. February 4(2): 33.

Introduction

Parasites pose a great risk to the animals. They cause huge economic losses to the livestock industry. It causes mortality of animals, reduces the fertility of animals and decreases the production of both the meat and milk of animals. There are different parasites which cause diseases in animals. These may be either ecto-parasites or endo-parasites [1]. In case of endo-parasitism various infections in ruminants like Fasciolosis, Schistosomiasis, Giardiasis, Amoebiasis, Ascariasis etc. are of major importance in livestock. In Pakistan, *Fasciola hepatica* is of major concern in the areas close to the canals, rivers or water bodies. It causes a disease named Fasciolosis in animals. It is a trematode and is commonly known as liver fluke [2].

Morphology

Liver fluke is a leaf shaped trematode. It measures about 3.5cm in length and 1.5cm in width. It is large enough to be seen with the naked eye. Its anterior portion is broader compared to the posterior portion. At the anterior end, a cone is present. Two suckers named oral and ventral are present [3].

Hosts

Fasciola hepatica affects a number of species of animals. The animals which may be infected include sheep, goats, cattle, buffalo, dog etc. Fasciolosis also affects man which means it is a zoonotic disease. Snail acts as intermediate host for the transmission of this disease [4].

Life Cycle

Fasciola hepatica lives in the bile duct. Its life cycle starts from eggs which are passed out in the feces. Hatching of these eggs takes place in the water and these develop into miracidium. It is the miracidium stage which infects the snail. In the snail, this miracidium undergoes further development into sporocyst, sporocyst into radia and radia converts into cercariae. Now these cercariae are released from the snail and attach to the vegetation. On this vegetation, these cercariae encyst to form the metacercariae. Metacercariae is the infective stage. These metacercariae are ingested by the definitive host and reach the small intestine. In the small intestine, these metacercariae excyst and reach the liver tissue by penetrating through the intestinal wall and peritoneal cavity. This further penetrates the liver tissue to the bile duct

References

- [1] Huntington B, Bernardo TM, Bondad-Reantaso M, Bruce M, Devleeschauwer B, Gilbert W, Grace D, Havelaar A, Herrero M, Marsh TL, Mesenowski S. Global Burden of Animal Diseases: a novel approach to understanding and managing disease in livestock and aquaculture. *Revue Scientifique Et Technique-Office International Des Epizooties*. 2021;40(2):567-83.
- [2] Regasa A, Seboka M. Review on Fasciolosis, its effect on meat quality/hazards and economical importances. *Entomol Ornithol Herpetol*. 2021;10:245.
- [3] Swarnakar G, Damor RN. Ultrastructural surface topography of *Fasciola gigantica* (trematoda: digenea) from infected liver of buffaloes (*Bubalus bubalis*) in Udaipur, India. *India. Pharmanest*. 2016;7(2):3056-9.
- [4] Mas-Coma S, Valero MA, Bargues MD. Human and animal fascioliasis: origins and worldwide evolving scenario. *Clinical microbiology reviews*. 2022 Dec 21;35(4):e00088-19.
- [5] Roberts EW. Studies on the life-cycle of *Fasciola hepatica* (Linnaeus) and of its snail host, *Limnaea* (*Galba*) *truncatula* (Müller), in the field and under controlled conditions in the laboratory. *Annals of Tropical Medicine & Parasitology*. 1950 Jul 1;44(2):187-206.
- [6] Fiss L, de Lourdes Adrien M, Marcolongo-Pereira C, Assis-Brasil ND, Sallis ES, Riet-Correa F, Ruas JL, Schild AL. Subacute and acute fasciolosis in sheep in southern Brazil. *Parasitology research*. 2013 Feb;112:883-7.

and become adult. These adult flukes now start laying eggs. These eggs travel to the duodenum with the bile and are passed out of the body of definitive host in feces [5].

Clinical Manifestation

Clinical manifestation of disease depends upon the number of metacercariae ingested. It may be either asymptomatic in cattle or may show severe clinical signs in sheep. It may be either acute or chronic. Acute form is seasonal while chronic form occurs in all seasons. In acute form, there is distended painful abdomen, anemia and sudden death. In chronic form, the signs observed include sub-mandibular edema, dullness, decreased milk production and anemia [6]. Fasciolosis also pre-disposes the animal to Black Disease which is caused by *Clostridium novyi* [7].

Diagnosis and Treatment

Diagnosis is based on history, clinical signs, post-mortem findings and detection tests like ELISA. Oval operculated eggs are also observed in feces. In post-mortem, immature worms in the parenchyma of liver, migratory tracts of worm and adult worms in the bile duct are observed. Plasma level of gamma-glutamyltransferase may also help in diagnosis as its level is increased in liver damage [8,9]. Various anthelmintics are used for the treatment of Fasciolosis in animals. The drug choice for the treatment of Fasciolosis is Triclabendazole as it is effective against all stage of the *Fasciola hepatica*. Other drugs which may be used are Albendazole, Oxylozanide, Nitroxylin and Closantel [10].

Control

Controlling this disease is thus very important to minimize the losses. Control is directed towards the overall herd health program. Limit the spread of disease by controlling population of snails. Treat the infected animals and do regular deworming of the animals. Drain off the land areas close to the shed of animals and spray molluskacides. Hence, we can avoid such parasitic losses in livestock industry by focusing on proper control measures, deworming at regular intervals, rotational grazing and proper hygiene [11].

- [7] El Shanawany EE, Fouad EA, Keshta HG, Hassan SE, Hegazi AG, Abdel-Rahman EH. Immunomodulatory effects of Moringa oleifera leaves aqueous extract in sheep naturally co-infected with *Fasciola gigantica* and *Clostridium novyi*. *Journal of parasitic diseases*. 2019 Dec;43:583-91.
- [8] Shahzad W, Mehmood K, Munir R, Aslam W, Ijaz M, Ahmad R, Khan MS, Sabir AJ. Prevalence and molecular diagnosis of *Fasciola hepatica* in sheep and goats in different districts of Punjab, Pakistan. *Pak Vet J*. 2012 Mar 11;32(4):535-8.
- [9] PILLAI AS. TRAFFIC MANAGEMENT: IMPLEMENTING AI TO OPTIMIZE TRAFFIC FLOW AND REDUCE CONGESTION. *Journal of Emerging Technologies and Innovative Research*. 2024 Jul 1;11(7).
- [10] Costa M, Saravia A, Ubios D, Lores P, da Costa V, Festari MF, Landeira M, Rodríguez-Zraquía SA, Bancho G, Freire T. Liver function markers and haematological dynamics during acute and chronic phases of experimental *Fasciola hepatica* infection in cattle treated with triclabendazole. *Experimental Parasitology*. 2022 Jul 1;238:108285.
- [11] Tabari MA, Vahdati SA, Samakkhah SA, Araghi A, Youssefi MR. Therapeutic efficacy of triclabendazole in comparison to combination of triclabendazole and levamisole in sheep naturally infected with *Fasciola* sp. *Journal of Parasitic Diseases*. 2022 Mar:1-7.