Biological Times Volume No. 4 | Issue No. 7 | Pages 48-49

The Deleterious Duo: Group A *Streptococcus* and *Staphylococcus* Aureus in Necrotizing Fasciitis

Muhammad Bilal Iftikhar*, Sarosh Zameer, Own Nasir, Nouman Tariq and Ahmad Raza

Faculty of Veterinary Science, University of Agriculture, Faisalabad, Pakistan

*Corresponding Author: **lalahadi2003@gmail.com**

ABSTRACT

Skin deterioration due to bacterial incursion is a symptomatic infection commonly known as Necrotizing fasciitis. The infection, being mono- or poly-microbial, is arduous to deal with. The disease is accelerating in the case of certain body manifestations. The microbes cause severe systemic toxicity through their virulence factors. The microbes adopt various mechanisms to bypass the body's natural defence system. Their laceration acts as a reservoir for pathogens that perforate the cutaneous barrier and become hostile to the body's defences. These noxious agents then enter various systemic routes and cause their malfunction. The malady is daunting to diagnose because the gangrenous tissue lacks pinpoint evidence of being caused by a specific infection. The LRINEC score also provides an exact tabulation of active malediction. Polymicrobial infestation is strenuous to confront, as antibiotics alone may not be effective and might entail surgical incision. The Assault by microbes makes the plight far more severe and Sisyphean to cope with. Proper vigilance and therapeutic practices should be deployed; otherwise, carnage can result throughout the globe. Debridement should be a prominent practice for wounds to prevent infection.

Keywords: mortification, monomicrobial, exotoxin, gangrenous tissue, virulence, agglomerating

To cite this article: Iftikhar MB, S Zameer, O Nasir, N Tariq & Ahmad Raza. The Deleterious Duo: Group A *Streptococcus* and *Staphylococcus* Aureus in Necrotizing Fasciitis. Biological Times. 2025. July 4(7): 48-49.

Introduction

Necrotizing fasciitis is used as a designation in order to cover a wide array of manifestations that occur in soft tissues of the body, especially the connective tissue fascia, causing soft tissue mortification [1]. The Deterioration is generally due to pre-existing underlying medical conditions; however, diabetes mellitus, immunosuppression, and chronic renal impairment are cited as principal co-morbidities with difficult prognostication in patients [2]. Tissue trauma often leads to the incursion of obnoxious pathogens, especially Group A Streptococcus (GAS), which is notoriously known for being the principal agent aggravating the conundrum. However, recently, there has been a great surge of occurrences that highlight the initiation of NF due to some unorthodox agents such as Staphylococcus aureus [3].

The dyad is considered to be one of the most noxious conventions during Fasciitis, as both of them can cause detrimental effects on the body. The Polymicrobial infection can cause acute septicemia and multiple organ dysfunction [4]. Necrotizing cellulitis and myositis are the major clinical markers that can be used to pinpoint the infection. Monomicrobial infection on itself is not that much menacing, although if the infection is accompanied by different strains of pathogen, it can be one of the most arduous to deal with [5]. Once it has impaired the cutaneous barrier, the control of infection is a daunting task, as we may not have to deal with only one but a tandem of microbes moving through the hematogenous route [6]. The nexus of GAS and S. aureus can destroy the body's immune system by their noxious mechanisms, either by releasing their virulence factors and certain hemotoxins [7].

What Causes their Virulence

GAS is a troublesome infection to treat as bacteria deploy numerous strategies to escape killing by the body's natural defense system, involving the evolution of antibiotic resistance, intracellular localization, and biofilm configuration. Due to the polymicrobial nature of the contamination, it becomes a reservoir for not only GAS but also for some other gram-negative species as S. aureus [8]. The agent being responsible for severe leukopenia and thrombocytopenia can collaterally cause equal damage to overall body functions. Both pathogens belong to the Type-II NF infection. GAS is a multifaceted microbe that targets the lymphoid and soft tissues of the body [9]. It propels severe inflammation employing different methodologies, such as inflammasomes, IL-1, and T-cell overdrive. Virulence factor of the microbe involves M proteins and Streptococcal pyogenic exotoxins; besides this, it causes skin mortification through Gasdermin mobilization [10]. The M protein, due to its hypervariable nature, assists it in dodging the body's natural defense mechanism through its N-terminal vicinity. The Pyogenic exotoxin B, being the most common virulence determinant, is a cysteine protease that arbitrates the bifurcation of transmembrane proteins, enables bacterial proliferation, and is the most common cause of Toxic shock syndrome (TSS), which is a major indication of Fasciitis. Capsule and DNase Sda-1 also help it avoid the body's defenses, either through cellular cohesion by capsule or through microflora synergism by DNase Sda-1 [11].

On the flip side, the microbial aggressors being produced by S. aureus are clumping factors A and B and fibronectin-binding protein (FBP). In the secretory toxins, toxic shock syndrome toxin-1 (TSST-1) and staphylococcal exfoliative toxin (SET) are the most prominent ones. The Clumping factor A acts as an FBP that aids the microbe in agglomerating to blood plasma, leading to septicemia, kidney abscess, infectious endocarditis, and several vascular invasions [12]. FBP is responsible for fibrin deactivation and disassembly. Toxin-1 occurrence is bound to bacterial emission of superantigen exotoxins, which are capable of initiating T cells through antigen-presenting cells. Through orthodox T cell triggering, antigen-presenting cells can engulf foreign bodies, process protease degradation and showcase them in a distinct binding spiral to the major histocompatibility complex class II, which, in turn, causes an immense expulsion of pro-inflammatory cytokines with notable manifestations as vascular discharge, arterial hypotensive mobility, organ dysfunction, and coagulation initiation [13]. Staphylococcal scalded skin disorder (SSSD) is one of the most prominent aftereffects noted by the intrusion and assault of exfoliative toxin, causing a fierce systemic retort. All these factors collectively contribute to the Pathogenicity of these microbes, thus also giving us a broad knowledge on how to deal with them in the case of obnoxious NF [14].

Diagnosis

Clinical diagnosis of the malady is an arduous task; commonly, the impairment due to GAS shows certain transitions in the body's natural functions, as there is localized inflammation with intense ache and ill-defined patterns of erythema. The Laboratory Risk Indicator for Necrotizing Fasciitis (LRINEC) score encompasses hemoglobin, glucose range, sodium, and creatinine waste. A tabulation greater than 6 suggests a high risk of NF. Another test, the Finger Stroke Test, implies that bifurcation of the cutaneous wound with far less deficiency signifies the outcome of NF due to GAS [15].

Conversely, S. aureus shows some extreme signs and symptoms as febrile fever, polypnea, and warm and inflamed appendages, although clinical tabulation might show C-reactive protein aggregate higher than 304 mg per liter (normal values are < 5 mg). Microbiological findings from the segmented part of gangrenous tissue, with the consistent use of vasopressor therapy, are the key diagnostic measurements that indicate septic shock syndrome. The ailment is the indicator for positive infection [16].

Antibiotic therapy during selective Necrotizing fasciitis:

Once the microbes have taken control of the body's natural responses and vitals, it's essential to input the right dose and medications to curtail their miasmic nature. As both the GAS and S. aureus belong to Type-II fasciitis infection. Therefore, penicillin, certain 2nd-generation cephalosporins, and macrolides are debated, although there are some speculations that clindamycin should be conjoined with beta-lactam drugs, but it is still vague regarding outcomes [17]. Infusion of broad-spectrum intravenous antibiotics is also significant as it can hamper the spread of infection. NF is undoubtedly a challenging complication to treat, as surgical incision and debridement are also needed to cope with such situations despite advanced bactericidal agents. The factor that makes the treatment a malediction is the manifestation of certain other ailments that are dwelling in the body. Diabetes mellitus and Anemia make the body cope with active infection, as they can lead to limb impairment and secondary tissue gangrene. Moreover, anemia can lead to a deficit of oxygen supply to the vital organs of the body, which in turn causes increased metabolic needs, leading to the shutdown of organs in no time. Bacterial invasion makes the case far more tormented and Sisyphean to deal with, thanks to proper restrictive measures and resources, which now enable us to confront the plight with a better understanding and intervention. Proper antibiotic treatment, surgical debridement, and bactericidal agents are the key control and preventive measures to successfully tackle further tissue festering, thereby preventing putrefaction. Disinfection of wounds should be part of medical interventions as it prevents further colonization from septic agents [18].

Conclusion

NF may not be the most formidable infection to have control over, but the proliferation of tandem either, unitarily or cooperatively annexed by certain metabolic or systemic instability, can be one of the most baneful and demise-causing infections. Proper medical vigilance and rehabilitation practices are imperative to note. Otherwise, extreme mortalities can result, and percolation of the disease can cause carnage throughout the world. Debridement and the care of wounds with therapeutics and antibiotics should be the pre-eminent treatment to prevent propagation of microbes.

References

[1] Tan JJ, Hui SJ, Tan SJ, Chan YH, Tham WY, Tan JY, Puhaindran ME, Hong CC. Necrotising fasciitis in the elderly: comparison between patients aged 60 years and older and those younger than 60 years. Singapore Medical Journal. 2025:10-4103.

[2] Nagayama J, Sato T, Takanori I, Kouji K, Mitsunobu N. Necrotising fasciitis with extensive necrosis caused by Lactobacillus: a case report. BMC Infectious Diseases. 2024 Apr 22;24(1):425.

[3] Allaw F, Wehbe S, Kanj SS. Necrotizing fasciitis: an update on epidemiology, diagnostic methods, and treatment. Current Opinion in Infectious Diseases. 2024 Apr 1;37(2):105-11.

[4] Becker DK, Röder M, Wagenhäuser J, Eichberger S, Herten M, Dudda M. Necrotizing fasciitis that led to unexpected and fulminant deterioration in less than 24 hours: a case report. Journal of International Medical Research. 2024 Dec;52(12):03000605241290480.

[5] Kochkine S, Payne DL, Chung K, Chen D, Bernstein MP, Baxter AB, McMenamy JM. Imaging of necrotizing fasciitis. Clinical Imaging. 2024 Dec 1;116:110331.

[6] Zhou L, Li H, Luo G. Expert Consensus on the Diagnosis and Treatment of Adult Necrotizing Fasciitis (2025 Edition). Burns & Trauma. 2025 Jun 12:tkaf031.

[7] Keller N, Boumasmoud M, Andreoni F, Tarnutzer A, von Matt M, Scheier TC, Epprecht J, Weller D, Gómez-Mejia A, Huemer M, von Reibnitz D. Investigating group A Streptococcus antibiotic tolerance in necrotizing fasciitis. Msphere. 2024 Sep 25;9(9):e00634-24.

[8] Pertea M, Luca S, Tatar R, Huzum B, Ciofu M, Poroch V, Palade DO, Vrinceanu D, Balan M, Grosu OM. Multidisciplinary Approach in Rare, Fulminant-Progressing, and Life-Threatening Facial Necrotizing Fasciitis. Infectious Disease Reports. 2024 Nov 1;16(6):1045-54.

[9] Nurmi K, Silventoinen K, Keskitalo S, Rajamäki K, Kouri VP, Kinnunen M, Jalil S, Maldonado R, Wartiovaara K, Nievas EI, Denita-Juarez SP. Truncating NFKB1 variants cause combined NLRP3 inflammasome activation and type I interferon signaling and predispose to necrotizing fasciitis. Cell Reports Medicine. 2024 Apr 16;5(4).

[10] Khan S, Ahmad R, Munir A, Nasir S, Adnan M, Naveed F, Idrees U, Fatima SM, Iqbal J. Trends in Necrotizing Fasciitis-Associated Mortality in the United States 2003–2020: A CDC WONDER Database Population-Based Study. World Journal of Surgery. 2025 May;49(5):1210-8.

[11] Deneubourg G, Schiavolin L, Lakhloufi D, Botquin G, Delforge V, Davies MR, Smeesters PR, Botteaux A. Nosocomial Transmission of Necrotizing Fasciitis: A Molecular Characterization of Group A Streptococcal DNases in Clinical Virulence. Microorganisms. 2024 Oct 31;12(11):2209.

[12] Mei M, Lu M, Li S, Ren X, Xing B, Hu Y, Wu Y, Chen H, Wang L, Yi L, Ming K. Development of nanobodies specific to clumping factors A of Staphylococcus aureus by yeast surface display. International Journal of Biological Macromolecules. 2024 Feb 1;259:129208.

[13] Touaitia R, Ibrahim NA, Abdullah Almuqri E, Basher NS, Idres T, Touati A. Toxic Shock Syndrome Toxin-1 (TSST-1) in Staphylococcus aureus: Prevalence, Molecular Mechanisms, and Public Health Implications. Toxins. 2025 Jun 24;17(7):323.

[14] Rouva G, Vergadi E, Krasagakis K, Galanakis E. Understanding host's response to staphylococcal scalded skin syndrome. Acta Paediatrica. 2025 Feb;114(2):241-7.

[15] Stallwood-Hall C, Binu V, Rajendran RK, Hiremath BV. Validating the laboratory risk indicator for necrotizing fasciitis (LRINEC) score: a prospective cohort study in a resource limited setting. ANZ Journal of Surgery. 2024 Mar;94(3):457-60.

[16] Plante TB, Juraschek SP, Howard G, Howard VJ, Tracy RP, Olson NC, Judd SE, Kamin Mukaz D, Zakai NA, Long DL, Cushman M. Cytokines, C-reactive protein, and risk of incident hypertension in the REGARDS study. Hypertension. 2024 Jun;81(6):1244-53.

[17] Rose W. Resurgence of group A streptococcal infections. Indian Journal of Pediatrics. 2025 Jun 13:1-7.

[18] Rao BS, Pavan M, Mutheeswaraiah Y, Rani GP, Babu PS, Mohan A, Jayaprada R, Sachan A, Lahari SS. A comparative study of necrotising fasciitis in patients with and without type 2 diabetes mellitus. Journal of Clinical and Scientific Research. 2024 Jul 1;13(3):170-7.