Biological Times Volume No. 4 | Issue No. 7 | Pages 74-75

Integrated Control of Ticks and Mites

Zain Iftikhar[1], Muhammad Mubashir Rafiq[2], Muhammad Abdullah Qureshi[2*] and Syed Rizwan Ahmad[2]

1. Department of Entomology, Faculty of Agriculture, University of Agriculture Faisalabad
2. Faculty of Veterinary Science, University of Agriculture Faisalabad

*Corresponding Author: **abdullah5902070@gmail.com**

ABSTRACT

Ticks and mites are economically significant ectoparasites that infest a wide range of domestic and wild animals, as well as humans. Their presence affects livestock productivity by causing direct irritation, skin damage, blood loss, and transmitting various pathogens. Traditional control methods primarily relied on chemical acaricides; however, the emergence of resistance, environmental contamination, and concerns about residue in animal products have highlighted the limitations of chemical control alone. Integrated control strategies represent a comprehensive, sustainable approach that combines chemical, biological, environmental, genetic, and management-based interventions. This article explores the biology and significance of ticks and mites, challenges in their control, and the key components of integrated tick and mite management, emphasizing the need for region-specific, species-specific, and economically viable programs to ensure long-term efficacy and ecological balance.

Keywords: Ticks, Mites, Integrated pest management, Acaricide resistance, Livestock health

To cite this article: Iftikhar Z, MM Rafiq, MA Qureshi & SR Ahmad. Integrated Control of Ticks and Mites. Biological Times. 2025. July 4(7): 74-75.

Introduction

Ticks and mites belong to the subclass Acari and are major arthropod ectoparasites of veterinary importance [1]. Ticks are obligate blood-feeding parasites, whereas mites can be parasitic or free-living and are associated with a range of dermatological, respiratory, and systemic diseases in animals and humans. In livestock, these ectoparasites are responsible not only for direct harm but also for acting as vectors of serious infectious agents, including viruses, bacteria, protozoa, and rickettsiae. For decades, control efforts have heavily relied on synthetic acaricides. While effective initially, these chemicals have led to the rapid development of resistance in tick and mite populations, threatening the sustainability of control programs. In addition to resistance, issues such as environmental pollution, non-target species toxicity, and consumer concerns regarding chemical residues have necessitated a shift toward more holistic and sustainable strategies. Integrated control approaches aim to address these concerns by combining multiple, complementary techniques to minimize parasite loads, slow resistance development, and maintain ecological integrity.

Integrated management of Ticks and Mites

The control of ticks and mites begins with a deep understanding of their biology, behavior, and life cycles [2]. Ticks undergo incomplete metamorphosis, progressing through egg, larva, nymph, and adult stages, with each stage requiring a blood meal in most species. They can be one-host, two-host, or three-host ticks, each presenting different challenges for control based on their host preferences and environmental persistence. Mites, on the other hand, exhibit a wide range of life histories and habitats. Parasitic mites such as *Psoroptes*, *Sarcoptes*, and *Demodex* infest the skin, causing mange and dermatitis, while others like *Dermanyssus* or *Ornithonyssus* infest poultry and rodents, causing severe irritation and stress. These differences highlight the importance of tailoring control strategies to the specific parasite species and local epidemiological conditions.

Chemical control remains an important component of integrated tick and mite management [3]. Acaricides, including organophosphates, pyrethroids, amidines, and macrocyclic lactones, are commonly used for treatment and prevention. However, improper application, underdosing, frequent use of the same class of chemicals, and lack of rotation have led to widespread acaricide resistance. To combat resistance, chemical treatments should be applied strategically, based on surveillance data and susceptibility testing. Acaricides should be used as part of a rotation plan to reduce the selection pressure on parasite populations. Spot treatments, targeted applications during high-risk periods, and the use of combination products can enhance efficacy while mitigating resistance development.

Biological control is another promising component of integrated strategies [4]. It involves the use of natural predators, pathogens, or antagonistic organisms to suppress tick and mite populations. Entomopathogenic fungi such as *Metarhizium anisopliae* and *Beauveria bassiana* have shown effectiveness against ticks under laboratory and field conditions. These fungi infect and kill ticks by penetrating their cuticle, thereby reducing their population over time. Similarly, predatory mites such as *Phytoseiulus persimilis* are effective in controlling pest mites on plants and may be adapted for veterinary use under controlled settings. Nematodes, parasitoid wasps, and bacteria are also under investigation as potential biocontrol agents. The challenge lies in optimizing their delivery, ensuring environmental compatibility, and integrating them effectively with other control methods.

Environmental management plays a critical role in tick and mite control, particularly in pasture-based and outdoor production systems. Habitat modification, such as clearing bushy areas, mowing tall grasses, and fencing off high-risk zones, can reduce the survival and proliferation of ticks. Livestock movement patterns can also be altered to avoid heavily infested areas during peak tick activity. Strategic grazing, rotational grazing, and controlled burns in certain ecosystems help limit tick populations by interrupting their life cycle and reducing host exposure. For mites, sanitation and hygiene are especially important. Regular cleaning and disinfection of animal housing, proper bedding management, and minimizing overcrowding reduce the environmental load of parasitic mites. In poultry production, systematic cleaning between flocks and the use of mite-proof materials in housing design are essential components of mite control.

Host resistance is another underutilized but promising area in integrated parasite control. Some livestock breeds exhibit natural resistance or tolerance to tick and mite infestations. For instance, *Bos indicus* cattle tend to be more tick-resistant than *Bos taurus* breeds, likely due to differences in skin thickness, grooming behavior, and immune response. Selective breeding for parasite resistance can reduce the need for chemical treatments and improve long-term sustainability. Marker-assisted selection and genomic tools are increasingly being applied to identify and propagate desirable traits in breeding programs. Moreover, enhancing the overall health and immunity of animals through proper nutrition, vaccination, and stress reduction indirectly improves their ability to withstand parasite infestations.

Vaccination represents a novel approach to tick control and has made significant strides in recent years. The development of anti-tick vaccines, such as those based on the *Bm86* antigen derived from the midgut of *Rhipicephalus microplus*, has shown moderate success in reducing tick burden and reproduction. Vaccinated animals develop antibodies that damage the tick's internal organs upon feeding, impairing survival and reproduction. While no commercial vaccines are yet available for mites, ongoing research into mite salivary proteins and allergens may pave the way for future developments. Vaccines have the advantage of avoiding resistance, being residue-free, and potentially conferring long-term protection with minimal environmental impact.

Education and community involvement are vital for the success of any integrated control program. Farmers, veterinarians, and animal handlers must be trained to recognize early signs of infestation, understand the biology of the parasites, and implement control measures correctly. Misuse of acaricides, inadequate treatment intervals, and poor compliance with control protocols can undermine the effectiveness of integrated approaches. Extension services, awareness campaigns, and participatory research can foster local ownership and ensure that control measures are culturally acceptable and economically feasible.

Monitoring and surveillance are essential components of integrated tick and mite control. Regular inspection of animals, acaricide resistance testing, and environmental assessments provide critical data for decision-making.

Geographic information systems (GIS), remote sensing, and predictive modeling help identify high-risk areas and forecast outbreaks. This information supports the timing of interventions and allocation of resources, making control programs more efficient and cost-effective.

Policy support and regulatory frameworks also play a role in facilitating integrated control. Governments and veterinary authorities must ensure the availability of quality acaricides, promote research into alternative control methods, and enforce regulations on drug usage and residue limits. Subsidizing vaccines or biologics, providing diagnostic services, and supporting breeding programs for resistant animals are important policy interventions that can enhance the adoption of integrated practices.

The integration of traditional knowledge with modern science adds another layer of depth to tick and mite control [5]. In many regions, farmers use ethnoveterinary remedies such as plant extracts, oils, and ashes to repel or kill ectoparasites. While scientific validation is still ongoing, some of these remedies—like neem oil, garlic extracts, and tobacco-based solutions—have shown acaricidal activity. Combining these practices with evidence-based methods can enhance acceptance and provide affordable options, particularly in resource-limited settings.

Conclusion

Integrated control of ticks and mites offers a sustainable, multidimensional approach to managing ectoparasite infestations in animals. By combining chemical treatments with biological agents, environmental management, host resistance, vaccination, and education, integrated strategies address both the immediate and long-term challenges posed by these pests. Unlike single-method approaches, integrated control reduces the risk of acaricide resistance, minimizes environmental and public health risks, and enhances the resilience of livestock production systems. Success in integrated tick and mite control depends on local adaptation, continuous monitoring, stakeholder engagement, and policy support. As research continues to advance and new tools become available, integrated approaches will remain the cornerstone of effective ectoparasite management in both animal and public health domains.

References

[1] Mandal SC. Veterinary Entomology and Acarology. InTextbook of Veterinary Parasitology 2025 Apr 1 (pp. 385-472). Singapore: Springer Nature Singapore.

[2] Rizwan HM, Naz S, Raza M, Iqbal A, Iftakhar T, Abbas H, Akhtar T. Biology and ecology of parasites. InParasitism and Parasitic Control in Animals: Strategies for the Developing World 2023 Jul 10 (pp. 1-20). GB: CABI.

[3] Makwarela TG, Seoraj-Pillai N, Nangammbi TC. Tick control strategies: Critical insights into chemical, biological, physical, and integrated approaches for effective hard tick management. Veterinary sciences. 2025 Feb 2;12(2):114.

[4] Jaiswal DK, Gawande SJ, Soumia PS, Krishna R, Vaishnav A, Ade AB. Biocontrol strategies: an eco-smart tool for integrated pest and diseases management. BMC microbiology. 2022 Dec 30;22(1):324.

[5] Angon PB, Mondal S, Jahan I, Datto M, Antu UB, Ayshi FJ, Islam MS. Integrated pest management (IPM) in agriculture and its role in maintaining ecological balance and biodiversity. Advances in Agriculture. 2023;2023(1):5546373.