Biological Times

Volume No. 4 | Issue No. 8 | Pages 15-16

Neuroplasticity and its Role in Learning and Recovery after Brain Injury

Aqsa Shahid and Hina Parvez

Department of Physiology, The University of Faisalabad, Pakistan

*Corresponding Author: **aqsashahid6565@gmail.com**

ABSTRACT

The study aimed to understand the role of neuroplasticity in learning and recovery following brain injury, with an emphasis on how brain reorganizes itself to make up for lost function. Neuroplasticity is brain's capacity to generate new neural connections and is essential for learning new skills as well as recovery from brain injury. The study covered various forms of neuroplasticity including synaptic, structural, and functional plasticity and how these processes develop the basis of cognitive flexibility and functional recovery. Findings indicated that neuroplasticity allows functional compensation after brain injury like stroke and traumatic brain injury, and synaptic plasticity was helped accountable for strengthening neural connections. Rehabilitations strategies, including physical exercises, neurostimulation, and cognitive therapies, were identified to enhance neuroplastic changes, optimizing recovery outcomes.

Keywords: Neuroplasticity, Brain Injury, Long-Term Potentiation (LTP)

To cite this article: Shahid A & H Parvez. Neuroplasticity and its Role in Learning and Recovery after Brain Injury. Biological Times. 2025. August 4(8): 15-16.

Introduction

Neuroplasticity is brains' ability to reorganise itself by developing new neural pathways is a processes providing the platform for learning and rehabilitation [1]. It holds a remarkable enabling the brain to alter itself to environmental stimuli, acquire new skills, and recover from trauma. Neuroplasticity is not only necessary during childhood when the brain, when the brain is highly plastic but also throughout life, as the platform for ongoing learning and cognitive flexibility [2]. After brain injuries, including strokes, traumatic brain injuries, or neurodegenerative disorders, neuroplasticity is crucial for recovery by helping the brain reconstruct itself, replacing lost functions [3.4]. Neuroplasticity is a basic process that enables brain to adapt to environmental changes and compensate for damage, thus an essential mechanism for learning and recovery [5, 6].

In the middle of neuroplasticity are two basic mechanisms that include synaptic plasticity and structural plasticity. Synaptic plasticity refers to the change of synaptic influences between neurons in terms of strength and efficacy [7]. As defined by [8], when two neurons are activated closely together, their synaptic connection will be stronger, a phenomenon known as Long-Term Potentiation (LTP). LTP is the learning and memory mechanism at the cellular level, because it makes the connectivity between neurons stronger so that the brain can process information more effectively [9]. On the other hand, when neurons are not activated simultaneously, their synapse fades away through long-term depression, so that the brain can strengthen and refine neural circuits in such a way that only beneficial connections become solidified but less beneficial ones are eliminated [10]. These synaptic changes are at the center of learning new skills, memory, and plasticity to environmental changes. Neuroplasticity is a very complex process, which is why it has a variety of forms of plasticity that affect organization and work in the brain. [11].

Structural neuroplasticity can be defined as the alterations in the physical structure of networks and neurons as well as the number, shape, the strength, and interconnectivity of synapses [12], through which the brain adapts to changing environments and experiences. Structural plasticity has been demonstrated to occur during development and to continue through adulthood [13, 14]. Functional neuroplasticity, however, describes changes in neural network properties that include efficiency, strength, and synchrony of synapse changes. Functional plasticity is rapid, impacting a multitude of cognitive and behavioural processes in attention, memory, and perception [6]. Structure and function of brain adaptability is illustrated in figure 1 describing how the brain makes and remakes connections. The brain builds new brain networks when people gain experience as reflected in structural changes in the left. When the brain is damaged some new neurons build up to replace lost functions while older ones degenerate or rewire to enable these new connections, speeding recovery and overall brain function.

In addition to synaptic changes, neuroplasticity also involves structural changes in the brain. These include the growth of dendrites, the neuronal extensions that take input from other cells, and neurogenesis, the creation of new neurons [15, 16]. Dendritic growth increases the number of synapses a neuron can make, allowing for more complex and efficient communication within neural circuits [17]. Neurogenesis most commonly occurs in areas like the hippocampus, an area crucial for learning and memory, and is crucial for cognitive flexibility [18]. In accordance with [19], such structural adaptation has special significance when the brain has been damaged, for instance, after stroke or Traumatic Brain Injury (TBI). For such instances, [12] argued that, neuroplasticity allowing the brain to rearrange itself into making new connections in replacement of the lost functions.



Figure 1: Diagram illustrating structural and functional neuroplasticity, showing how the brain reorganizes and forms new connections following learning, experience, and injury recovery.

The Role of Neuroplasticity in Learning

Neuroplasticity is an imperative aspect of learning as it allows the brain to change, rearrange, and reconnect new neural networks during experience, challenge, and developing new skills [20]. At the root of neuroplasticity is synaptic plasticity, which entails the strengthening or weakening of neural connections among neurons. As per [21], this is achieved by mechanisms such as LTP and LTD wherein repeated stimulation of neurons makes their communication more effective. Learning activities whether through studying, practicing a skill, or simply interacting with the environment have been found to result in alteration in the brain's structure and function [22]. For instance, it has been demonstrated that when individuals acquire a new task, such as a new language or musical instrument, the brain regions involved in those tasks alter rather significantly structurally [23, 24]. This can be observed in heightened dendritic growth, synaptic remodeling, and even neurogenesis, particularly in regions such as the hippocampus, and prefrontal cortex, which normal individuals have to do with memory, attention, and execution. A study discovered that difficult learning tasks such as playing music or learning a new language stimulate neuroplasticity by raising gray matter in parts of the brain involved with sensory-motor functions and memory [25]. This is the way the brain streamlines and reorganizes its circuitry to save novel information, which is the basis for learning. Moreover, practice and repetition reinforce acquired information and strengthen the neural pathways responsible for its retention.

The principle of "use it or lose it" applies, with connections that are not routinely stimulated growing weaker and eventually being pruned off [26]. While neuroplasticity is most pronounced in childhood, adults nonetheless benefit from it, albeit at a slower rate. Mental stimulation and physical activity can still cause significant neuroplastic changes in adults [1]. However, neuroplasticity also creates maladaptive changes, such as reinforcing negative neural circuits for addiction, depression, or PTSD [10]. Therefore, the shaping of neuroplasticity with structured environments and

therapeutic interventions is crucial. In education and cognitive rehabilitation, neuroplasticity is increasingly being used to develop effective learning strategies and therapies [27].

Neuroplasticity in Brain Injury Recovery

Brain injury through any source is an immediate concern for all physicians and patients with such condition [12]. Medical tradition in the past had to contend with brain injuries as irreversible conditions with very little prospect of recoveries or cures. Following a TBI or stroke, neuroplasticity is initiated, initiating a cascade of mechanisms designed to remap damaged regions of the brain [28]. This repair begins with secondary neuronal networks activation that substitute damaged areas, followed by synaptogenesis and neurogenesis, two activities that contribute to the reconstruction and recovery of brain function [12]. It has been reported that, neuroplasticity is an ongoing dynamic process even in adulthood, enabling the brain to form new connections as a result of damage as well as rehabilitation [29]. The result of neuroplasticity in recovery depends heavily on the intensity and timing of rehabilitative therapy.

Initiation of early rehabilitation following injury also maximizes neuroplastic potential by stimulating the brain during the most plastic period [28]. Ideal therapies, e.g., physical therapy maneuvers, neurostimulation, and pharmacological interventions, play critical roles in enhancing neuroplastic modifications. Transcranial Magnetic Stimulation (TMS), for instance, has been shown to hold promising outcomes in stimulating parts of the brain to restore motor and cognitive function by modulating cortical plasticity [30]. Also, the presence of Brain-Derived Neurotrophic Factor (BDNF) has been strongly implicated in neuroplastic recovery because BDNF is involved in supporting neuronal survival, growth, and reorganization of synapses, making it possible for recovery in the affected brain areas (31). Even though younger patients will tend to have more neuroplasticity, existing studies suggest that even old adults and severely injured patients can be treated by certain neuroplastic therapies [28].



Figure 2: Schematic representation of neuroplasticity at different levels.

Following a brain injury, neuroplasticity can be adaptive or maladaptive. Adaptive neuroplasticity refers to compensation in function through rerouting of neuronal circuits, but maladaptive plasticity potentially worsens recovery if compensatory adjustments can hinder healing of the primary neural circuits [32]. Acute plasticity changes, such as synaptic strengthening and neurotransmitter release, arise rapidly after injury, whereas late changes, such as axonal sprouting and dendritic reorganization, have the potential to facilitate long-term recovery. Effective rehabilitation methods leverage neuroplasticity to promote recovery. Rehabilitation techniques like Constraint-Induced Movement Therapy (CIMT), Repetitive Task Training (RTT), and cognitive rehabilitation use experience-dependent plasticity to amplify neural circuits [28]. Furthermore, emerging technology in the guise of Virtual Reality (VR), Brain-Computer Interfaces (BCIs), and noninvasive brain stimulation like Transcranial Magnetic Stimulation (TMS) are being formulated as effective neurorehabilitation tools, offering individually tailored pathways to recovery [33]. These treatments trigger motor and cognitive recovery by stimulating brain regions and promoting plasticity. Furthermore, individualised neurorehabilitation where treatments are customised to patient profiles has proven to be successful strategy for maximising recovery outcomes [34].

Conclusion

The study revealed that, neuroplasticity is a driving force in learning and recovery from brain injury. It observed that the ability of brain to reorganise and create new neural pathways helped compensation of lost functions after trauma such as strokes and traumatic brain injurers. The brain compensated by strengthening or weakening synaptic contacts and reforming neutral networks through processes like synaptic plasticity and structural plasticity. It further highlighted that, accurate rehabilitation methods, including physical training, neurostimulation and cognitive rehabilitation enhanced neuroplastic changes, thereby enhancing recovery outcomes. Additionally the study highlighted the need of early intervention and the need of individualized rehabilitation programs based on individual needs and increased the neuroplastic capacity of the brain and led to functional recovery in the patient.

References

[1] Puderbaugh M, Emmady PD. Neuroplasticity. InStatPearls [Internet] 2023 May 1. StatPearls Publishing.
[2] Chen SA, Goodwill AM. Neuroplasticity and adult learning. InThird international handbook of lifelong learning 2022 Sep 10 (pp. 1-19). Cham: Springer International Publishing.
[3] Rajan RK. A comprehensive review on adaptive plasticity and recovery mechanisms post-acquired brain injury. Neuroprotection. 2025.
[4] Su S, Veeravagu A, Grant G. Neuroplasticity after traumatic brain injury.
[5] Voss P, Thomas ME, Cisneros-Franco JM, de Villers-Sidani É. Dynamic brains and the changing rules of neuroplasticity: implications for learning and recovery. Frontiers in psychology. 2017 Oct 4;8:274878.
[6] Marzola P, Melzer T, Pavesi E, Gil-Mohapel J, Brocardo PS. Exploring the role of neuroplasticity in development, aging, and neurodegeneration. Brain sciences. 2023 Nov 21;13(12):1610.
[7] Kania BF, Wrońska D, Zięba D. Introduction to neural plasticity mechanism. Journal of Behavioral and Brain Science. 2017 Feb 6;7(2):41-9.
[8] Bin Ibrahim MZ, Benoy A, Sajikumar S. Long-term plasticity in the hippocampus: maintaining within and 'tagging'between synapses. The FEBS journal. 2022 Apr;289(8):2176-201.
[9] Caya-Bissonnette L, Béïque JC. Half a century legacy of long-term potentiation. Current Biology. 2024 Jul 8;34(13):R640-62.
[10] Gazerani P. The neuroplastic brain: current breakthroughs and emerging frontiers. Brain Research. 2025 Apr 23:149643.
[11] Frizzell TO, Phull E, Khan M, Song X, Grajauskas LA, Gawryluk J, D'Arcy RC. Imaging functional neuroplasticity in human white matter tracts. Brain Structure and Function. 2022 Jan;227(1):381-92.
[12] Joshua AM. Neuroplasticity. InPhysiotherapy for adult neurological conditions 2022 Jun 21 (pp. 1-30). Singapore: Springer Nature Singapore.
[13] Chen X, Zhou X, Yang L, Miao X, Lu DH, Yang XY, Zhou ZB, Kang WB, Chen KY, Zhou LH, Feng X. Neonatal exposure to low-dose (1.2%) sevoflurane increases rats' hippocampal neurogenesis and synaptic plasticity in later life. Neurotoxicity research. 2018 Aug;34:188-97.
[14] Garthe A, Roeder I, Kempermann G. Mice in an enriched environment learn more flexibly because of adult hippocampal neurogenesis. Hippocampus. 2016 Feb;26(2):261-71.
[15] Kania BF, Wrońska D, Zięba D. Introduction to neural plasticity mechanism. Journal of Behavioral and Brain Science. 2017 Feb 6;7(2):41-9.
[16] Tataranu LG, Rizea RE. Neuroplasticity and Nervous System Recovery: Cellular Mechanisms, Therapeutic Advances, and Future Prospects. Brain Sciences. 2025 Apr 15;15(4):400.
[17] Drigas AS, Karyotaki M, Skianis C. An integrated approach to neuro-development, neuroplasticity and cognitive improvement. International Journal of Recent Contributions from Engineering, Science & IT (iJES). 2018 Nov;6(3):4-18.
[18] Anacker C, Hen R. Adult hippocampal neurogenesis and cognitive flexibility—linking memory and mood. Nature Reviews Neuroscience. 2017 Jun;18(6):335-46.
[19] Ngwenya LB, Danzer SC. Impact of traumatic brain injury on neurogenesis. Frontiers in neuroscience. 2019 Jan 9;12:1014.
[20] Xie L. Neuroplasticity the Brain's Remarkable Ability to Adapt and Change. Health Science Journal. 2024;18(4):1-3.
[21] Maffei A. Long-term potentiation and long-term depression. InOxford research encyclopedia of neuroscience 2018 Feb 26.
[22] McCandliss B, Toomarian E. Putting Neuroscience in the Classroom: How the Brain Changes As We Learn [Internet]. Pew.org. The Pew Charitable Trusts; 2020 [cited 2025 Jul 6]. Available from: https://www.pew.org/en/trend/archive/spring-2020/putting-neuroscience-in-the-classroom-how-the-brain-changes-as-we-learn
[23] Wei X, Gunter TC, Adamson H, Schwendemann M, Friederici AD, Goucha T, Anwander A. White matter plasticity during second language learning within and across hemispheres. Proceedings of the National Academy of Sciences. 2024 Jan 9;121(2):e2306286121.
[24] Olszewska AM, Gaca M, Herman AM, Jednoróg K, Marchewka A. How musical training shapes the adult brain: Predispositions and neuroplasticity. Frontiers in neuroscience. 2021 Mar 10;15:630829.
[25] Kwatra, M. and Ahuja, A., 2025. Reshaping the Brain Through Music: Instrumental Learning, Neuroplasticity, and Academic Achievement. Sangeet Galaxy, 14(1).
[26] Kukreja J, Manoharan G. Rewiring the Mind: Unlocking the Healing Power of Neuroplasticity. InAdvancing Medical Research Through Neuroscience 2025 (pp. 239-256). IGI Global Scientific Publishing.
[27] Kumar J, Patel T, Sugandh F, Dev J, Kumar U, Adeeb M, Kachhadia MP, Puri P, Prachi FN, Zaman MU, Kumar S. Innovative approaches and therapies to enhance neuroplasticity and promote recovery in patients with neurological disorders: a narrative review. Cureus. 2023 Jul 15;15(7).
[28] Zotey V, Andhale A, Shegekar T, Juganavar A. Adaptive neuroplasticity in brain injury recovery: strategies and insights. Cureus. 2023 Sep 24;15(9).
[29] Tataranu, L.G. and Rizea, R.E., 2025. Neuroplasticity and Nervous System Recovery: Cellular Mechanisms, Therapeutic Advances, and Future Prospects. Brain Sciences, 15(4), p.400.
[30] Sharbafshaaer M, Cirillo G, Esposito F, Tedeschi G, Trojsi F. Harnessing Brain Plasticity: The Therapeutic Power of Repetitive Transcranial Magnetic Stimulation (rTMS) and Theta Burst Stimulation (TBS) in Neurotransmitter Modulation, Receptor Dynamics, and Neuroimaging for Neurological Innovations. Biomedicines. 2024 Nov 1;12(11):2506.
[31] Bazzari AH, Bazzari FH. BDNF therapeutic mechanisms in neuropsychiatric disorders. International journal of molecular sciences. 2022 Jul 29;23(15):8417.
[32] Rajan, R.K., 2025. A comprehensive review on adaptive plasticity and recovery mechanisms post-acquired brain injury. Neuroprotection.
[33] Reddy KJ. Neuroplasticity and Technology. InInnovations in Neurocognitive Rehabilitation: Harnessing Technology for Effective Therapy 2025 Apr 27 (pp. 137-169). Cham: Springer Nature Switzerland.
[34] Du W, Shen J, Su T. Neuroplasticity in Stroke Rehabilitation: Harnessing Brain's Adaptive Capacities for Enhanced Recovery. Journal of Innovations in Medical Research. 2023 Nov 30;2(11):50-8.